

PAINEL 5

VARIATIONS OF THE SOLAR DIAMETER

Eugênio Reis Neto¹, **Alexandre Humberto Andrei**^{1,2},
Jucira Lousada Penna¹, **Evgueni Jilinski**^{1,3,4}, **Victor d'Ávila**¹
 1 - ON/MCT
 2 - GEA/OV/UFRJ
 3 - LNCC/MCT
 4 - Pulkova Observatory

This work aims to present the final results of the 1998 to 2000 campaign of solar diameter surveying. The employed instrument was a Danjon astrolabe, at the Observatório Nacional campus, and specially modified for the solar observations. During the time lapse, 10807 independent measurements of the solar diameter were made, Eastwards and Westwards from the local meridian and evenly distributed. A study is made to identify the systematic effects of the observational conditions upon the reduction final outcome. The mean temperature at the the moment of the observation is shown as the most influential parameter upon the final result. Next to it, follows the temperature variation, the Fried's factor, and the standard deviation of the reflected parabola, all these presenting a minor and complex degree of influence. The derived corrections are of the order of hundredths of arc seconds, thus being tenfold smaller than the typical error of one observation. The mean semi-diameter for the time lapse is $959''.107 \pm 0''.006$. Through the use of a CLEAN algorithm the periodic terms of the semi-diameter are obtained. The largest amplitude is attached to that of 515 days. By using a second type of algorithm, namely the DCDFIT, the found periods stand additional proof. A dependency of the semi-diameter on the observed heliolatitude is verified. The difference between the equatorial and polar radii calculated is $\Delta r = 0''.013 \pm 0''.004$, and the solar quadrupole moment inferred is $|J_2| = (3.61 \pm 2.90) \times 10^{-6}$.



COSMOLOGIA

PAINEL 6

BRANE WORLD COSMOLOGIES AND STATISTICAL PROPERTIES OF GRAVITATIONAL LENSES

J. S. Alcaniz¹, **D. Jain**², and **A. Dev**²
 1 - Astronomy Department, University of Washington, Seattle,
 Washington, 98195-1580, USA
 2 - Dept. of Physics and Astrophysics, University of Delhi, Delhi-110007, India

Brane world cosmologies seem to provide an alternative explanation for the present accelerated stage of the Universe with no need to invoke either a cosmological constant or an exotic *quintessence* component. Here we investigate statistical properties of gravitational lenses for some particular scenarios based on this large scale modification of gravity. We show that a large class of such models are compatible with the current lensing data for values of the matter density parameter $\Omega_m \leq 0.94$ (1σ). If one fixes Ω_m to be ≈ 0.3 , as suggested by most of the dynamical estimates of the quantity of matter in the Universe, the predicted number of lensed quasars requires a slightly open universe with a crossover distance between the 4 and 5-dimensional gravities of the order of $1.76 H_0^{-1}$.

PAINEL 7

THE CORRELATION FUNCTION BETWEEN HIGH DENSITY PEAKS IN A MIXED DISTRIBUTION FLUCTUATION FIELD

A. P. A. Andrade¹, **A. L. B. Ribeiro**², **C. A. Wuensche**¹
 1 - Divisão de Astrofísica, INPE
 2 - Departamento de Matemática Aplicada, IMECC, Unicamp

We calculate the correlation function of high density peaks in the primordial fluctuation field from the probability density function of a mixed random field. The correlation function is estimated using an expansion of the two-dimensional probability density in a finite series of generalized Hermite polynomials, taking the quasi-moment functions as the expansion coefficients. We compare the estimated correlation functions for a Gaussian probability density field with a

general random non-Gaussian field in order to quantify small deviations from Gaussianity. Our main assumption is that the fluctuation field has a mixed probability density function (PDF) of the form: $P(\delta) = (1-\alpha)f_1(\delta) + \alpha f_2(\delta)$, where: a) f_1 is the probability distribution of an adiabatic Gaussian field, b) f_2 is the probability distribution of a general random non-Gaussian field and c) α is a mixing parameter which allows us to modulate the contribution of each component to the resultant field. From the above PDF, we derive the correlation function between high density peaks for small deviations from Gaussianity. We show how this mixed correlation function can be applied to Cosmic Microwave Background data to search for possible deviations of a Gaussian signature in the statistical properties of the temperature fluctuations.

PAINEL 8

GAMMA-RAY BURSTS DEMOGRAPHY FROM THE SAMPLE WITH KNOWN REDSHIFTS

César A. Costa, Francisco Jablonski
INPE/MCT

Gamma-ray Bursts (GRB) are the most energetic events in nature on short time-scales, being second only to the Big-Bang itself. The mechanism for generating the enormous radiant energy output (with isotropic peak luminosities in excess of 10^{52} erg/s) is not fully understood, but there may be a connection with supernova events. The number of GRB with known redshifts is now large enough for statistical studies to be carried out. In this contribution we examine the constraints that can be put on the luminosity function of GRB and their space distribution by examining the predictions of simple models and comparing them to the observed redshift distribution, the observed $\log N \times \log S$ diagram and the observed distribution of luminosities. Preliminarily, we have verified that strong evolution effects are needed to explain the redshift distribution, both for the single luminosity case (as if GRB were standard candles) or spread luminosity functions. The evolution term has the form $(1+z)^\alpha$ with $\alpha \approx 3.5$. One key point in our analysis is that we chose the smoothest possible functionals to describe both the luminosity function and space distribution. This is important since there is a clear evidence for a bimodal distribution of peak luminosities. This work was supported by FAPESP 01/14527-3.

PAINEL 9

DEFLATIONARY COSMOLOGY: CONSTRAINTS FROM ANGULAR SIZE AND AGES OF GLOBULAR CLUSTERS

J. V. Cunha¹, J. S. Alcaniz², J. A. S. Lima¹
1 - DFTE/UFRN
2 - Astronomy Department/UW

Observational constraints to a large class of decaying vacuum cosmologies are derived using the angular size data of compact radio-sources and the latest age estimates of globular clusters. For this class of deflationary $\Lambda(t)$ models, the present value of the vacuum energy density is quantified by a positive β parameter smaller than unity. In the case of milliarsecond compact radio-sources, we find that the allowed intervals for β and the matter density parameter Ω_m are heavily dependent on the value of the mean projected linear size l . Constraints from age estimates of globular clusters and old high redshift galaxies are not so restrictive, thereby suggesting that there is no age crisis for this kind of $\Lambda(t)$ cosmologies.

PAINEL 10

A TIME VARYING SPEED OF LIGHT FOR A MODIFIED VACUUM

C. S. da Câmara Neto, J. C. Carvalho, M. R. de Garcia Maia, R. B. do Nascimento
UFRN

In the last few years, several cosmologies with time variation of the speed of light, c , have been proposed. In these schemes, the functional dependence of c with time is either imposed or obtained through the use of scalar fields (analogous to what happens with G in the Brans-Dicke theory). A consequence of such approach is that the form of the equations of Electrodynamics must be altered. In the present work, we derive a form for the time variation of c for a vacuum with time dependent electric permittivity and magnetic permeability, whereas requiring the validity of Maxwell Electrodynamics. We show that some forms for the time variation of c , that have been postulated in the literature, may be justified in our approach. By imposing some symmetry conditions on the electromagnetic wave equations, the time dependence of c is determined. The solution for these wave equations are derived and compared with the classical results.