

Adaptive multiresolution methods for evolutive PDEs

Margarete Oliveira Domingues

Laboratório Associado de Computação e Matemática Aplicada (LAC/CTE),
Instituto Nacional de Pesquisas Espaciais (INPE),
Av. dos Astronautas 1758, 12227-010 São José dos Campos, Brazil
E-mail: mo.domingues@lac.inpe.br

Sônia Maria Gomes

Universidade Estadual de Campinas, IMECC,
Caixa Postal 6065, 13083-970 Campinas SP, Brazil
E-mail: soniag@ime.unicamp.br

Olivier Roussel

Institut für Technische Chemie und Polymerchemie (TCP),
Universität Karlsruhe, Kaiserstr. 12, 76128 Karlsruhe, German
E-mail: roussel@ict.uni-karlsruhe.de

Kai Schneider

Laboratoire de Modélisation et Simulation Numérique en Mécanique et Génie des Procédés (MSNM-GP),
CNRS and Université d'Aix-Marseille,
38 rue Frédéric Joliot-Curie, 13451 Marseille Cedex 20, France,
Centre de Mathématiques et d'Informatique (CMI), Université de Provence,
39 rue Frédéric Joliot-Curie 13453 Marseille Cedex 13, France
E-mail: kschneid@cmi.univ-mrs.fr

ABSTRACT

Adaptive multiresolution strategies in space [1] and time [2] allow considerable speedup of second-order finite volume schemes for multi-dimensional evolutive partial differential equation in Cartesian geometry, while controlling the accuracy of the discretization. This strategy is based in a multiresolution technique for finite volume schemes with explicit time discretization using a local scale-dependent time stepping with control techniques like Embedded Runge-Kutta [?]. An adaptive grid is introduced by suitable thresholding of the wavelet coefficients, which maintains the accuracy of the finite volume scheme of the regular grid. On the finest scale the size of the time step is imposed by the stability condition of the explicit scheme. On larger scales, the time step can be increased without violating the stability requirement of the explicit scheme. The implementation uses a dynamic tree data structure. The accuracy and efficiency of this fully adaptive method is illustrated with new applications in one, two and three space dimensions

for some test cases illustrating the additional speed-up of these technique compared to both multiresolution scheme with global time stepping and finite volume scheme on a regular grid.

Referências

- [1] Roussel, O. and Schneider, K. and Tsigulin, A. and Bockhorn, H., A conservative fully adaptive multiresolution algorithm for parabolic PDEs, *J. Comput. Phys.*, 188 (2003), 493-523.
- [2] Domingues, M. O. and Gomes, S. M. and Roussel, O. and Schneider, K., An adaptive multiresolution scheme with local time-stepping for evolutionary PDEs, *J. Comput. Phys.*, 227 (2008), 3758-3780.
- [3] Space-time adaptive multiresolution methods for hyperbolic conservation laws: applications to compressible Euler equations, *Appl. Num. Math.* (in revision).