


1. Publication Nº <i>INPE-4527-PRE/1277</i>	2. Version	3. Date <i>April 1988</i>	5. Distribution <input type="checkbox"/> Internal <input checked="" type="checkbox"/> External <input type="checkbox"/> Restricted
4. Origin <i>DAS</i>	Program <i>ASTRO</i>		
6. Key words - selected by the author(s) <i>REDSHIFTS OF GALAXIES, CLUSTERS OF GALAXIES.</i>			
7. U.D.C.: <i>524.77</i>			
8. Title <i>NEW MEASUREMENTS OF RADIAL VELOCITIES IN CLUSTERS OF GALAXIES</i>		10. Nº of pages: <i>6</i>	
		11. Last page: <i>5</i>	
		12. Revised by	
9. Authorship <i>D. Proust¹ A. Talavera² E. Salvador-Solé³ A. Mazure¹ H.V. Capelato</i>		13. Authorized by <i>Marcos Antônio Raupp</i> <i>General Director</i>	
		Responsible author 	
14. Abstract/Notes <i>We have obtained 101 new redshifts for 100 galaxies. Data for individual galaxies are presented and the accuracy of velocities determined using four different instrumentations (located at Calar Alto, La Palma, ESO and Hawaii) is discussed.</i> ¹ <i>Observatoire de Paris, Section de Meudon et Université Paris VII, Département d'Astrophysique Fondamentale, F-92195 Meudon Principal Cedex, France.</i> ² <i>I.U.E. Observatory, European Space Agency, P.O. Box 54065, Madrid, Spain.</i> ³ <i>Departamento de Física de la Tierra y del Cosmos, Universidad Central de Barcelona, Av. Diagonal 647, Barcelona, 28, Spain.</i>			
15. Remarks <i>Published in Astron. Astrophys. Suppl. Ser., 67, 57-61, 1987.</i>			



PROPOSTA PARA
PUBLICAÇÃO

- ☐ DISSERTAÇÃO
☐ TESE
☐ RELATÓRIO
☒ OUTROS

TÍTULO

"NEW MEASUREMENTS OF RADIAL VELOCITIES IN CLUSTERS OF GALAXIES"

IDENTIFICAÇÃO

AUTOR(ES)

D. Proust
A. Talavera
E. Salvador Solé
A. Mazure
H.V. Capelato

ORIENTADOR

CO-ORIENTADOR

DIVULGAÇÃO

☒ EXTERNA ☐ INTERNA ☐ RESTRITA

EVENTO/MEIO

☐ CONGRESSO ☒ REVISTA ☐ OUTROS

LIMITE

DEFESA

CURSO

ORGÃO

DAS

___/___/___

___/___/___

___/___/___

REV. TÉCNICA

NOME DO REVISOR

RECEBIDO

DEVOLVIDO

ASSINATURA

___/___/___

___/___/___

___/___/___

NOME DO RESPONSÁVEL

Hugo Vicente Capelato

APROVADO

DATA

ASSINATURA

☒ SIM

☐ NÃO

___/___/___

___/___/___

REV. LINGUAGEM

Nº

PRIOR.

RECEBIDO

NOME DO REVISOR

___/___/___

___/___/___

___/___/___

PÁG.

DEVOLVIDO

ASSINATURA

___/___/___

___/___/___

___/___/___

OS AUTORES DEVEM MENCIONAR NO VERSO INSTRUÇÕES ESPECÍFICAS, ANEXANDO NORMAS, SE HOUVER

RECEBIDO

DEVOLVIDO

NOME DA DATILÓGRAFA

___/___/___

___/___/___

Neide Gêa

Nº DA PUBLICAÇÃO:

PÁG.:

CÓPIAS:

Nº DISCO:

LOCAL:

☐ SIM

☐ NÃO

___/___/___

AUTORIZO A PUBLICAÇÃO

DISS. OU TESE

APROVAÇÃO

DATILOGRAFIA

DIRETOR

OBSERVAÇÕES E NOTAS

Artigo publicado, porém sem publicação interna.

New measurements of radial velocities in clusters of galaxies (*)

D. Proust ⁽¹⁾, A. Talavera ⁽²⁾, E. Salvador Solé ⁽³⁾, A. Mazure ⁽¹⁾ and H. V. Capelato ⁽⁴⁾

⁽¹⁾ Observatoire de Paris, Section de Meudon et Université Paris VII, Département d'Astrophysique Fondamentale, F-92195 Meudon Principal Cedex, France

⁽²⁾ I.U.E. Observatory, European Space Agency, P.O. Box 54065, Madrid, Spain

⁽³⁾ Departamento de Física de la Tierra y del Cosmos, Universidad Central de Barcelona, Av. Diagonal 647, Barcelona 28, Spain

⁽⁴⁾ IAG-USP, Caixa Postal 30627, 01051 São Paulo, Brazil

Received December 12, 1985 ; accepted February 24, 1986

Summary. — We have obtained 101 new redshifts for 100 galaxies in ten clusters of galaxies. Data for individual galaxies are presented and the accuracy of velocities determined using four different instrumentations (located at Calar Alto, La Palma, ESO and Hawaii) is discussed.

Key words : redshifts of galaxies — clusters of galaxies.

1. Introduction.

From recent developments of observational astronomy, the overall view of the structure of the universe appears to be very different from the homogeneous and isotropic one claimed by the traditional cosmology. The scheme of long, interconnected linear filaments (of size $\approx 100 h^{-1}$ Mpc, where $h = H_0/100$ and H_0 is the Hubble constant) formed by concentration of galaxies seems now well established (e.g. Fontanelli, 1984), these regions being separated by large voids empty of bright galaxies.

One of the fundamental aspects of the understanding of such large structures is to accede to the third dimension in order to modify their « projected view ». For instance, the spatial correlation function $\xi(r)$ of galaxy pairs (Dekel, 1984) estimated from the Center for Astrophysics (CFA) survey shows unobserved structures from estimates made on the basis of catalogues which use only projected positions. The use of percolation applied to a sample of three-dimensional galaxies is also a powerful way to explain the reality of large structures.

If the redshift determination is the only way to have access to the third dimension, it is also essential for the understanding of the dynamics because it leads to the

determination of the velocity dispersion in particle systems. The cosmic virial theorem (Peebles, 1980) using two and three points spatial correlation functions gives estimates of the cosmological parameter Ω_0 .

The radial velocities are also essential for the understanding of structures like clusters of galaxies, their dynamical analysis leading to their mass and to an estimate of the missing mass in the universe.

Analyses of some Abell clusters have been recently published : Gregory and Thompson (1984) for A 2197 and A 2199 ; Geller *et al.* (1984) for A 1142 ; Mazure *et al.* (1985) for A 496, and in this paper, we present 101 new redshifts for 100 galaxies in ten clusters. The data for individual galaxies are presented and the accuracy of velocities determination using four different instrumentations is discussed.

2. Observations.

The program of radial velocity measurements was carried out between March 1984 and March 1985, i) at the 2.2 m telescope at Calar Alto (Spain), ii) at the 2.5 m Isaac Newton telescope at La Palma (Canary Islands) and iii) at the 1.50 m telescope at ESO (Chile). We also include results of observations made during a technical run at the 3.60 m Canada-France-Hawaii (CFH) telescope at Mauna Kea.

2.1 OBSERVATIONS AT CALAR ALTO. — Observations were made in two observing runs of six nights each with the 2.20 m telescope in March 1984 and March 1985.

(*) Based on observations made at Calar-Alto Observatory (Spain), La Palma Observatory (Canary Islands), European Southern Observatory — La Silla (Chile) and Canada — France — Hawaii Telescope operated by the National Research Council of Canada, the Centre National de la Recherche Scientifique of France, and the University of Hawaii.

Fifty per cent of the nights were suitable for observations because of weather conditions and Moon light.

We used the Boller & Chivens Spectrograph with an effective dispersion of 89 Å/mm. The spectral region covered was from 3900 Å to 7200 Å with a 600 line/mm grating blazed at 5500 Å. The slit was 180 µm widened, the resolution was about 8 Å FWHM. A two-stage EM9916 image intensifier of 40 mm cathode diameter was used, cooled at -18°. Spectrograms were obtained on 11a0 Kodak plates and the wavelength calibration was made using a Fe-Ne Source before and after each spectrum. Exposure times used ranged between 1800 and 3600 s, depending on magnitude and weather conditions. The resulting signal to noise ratio was between 8 and 12.

Spectrograms were recorded with the PDS1010G microphotometer of the Institut d'Optique at Orsay (using a 20 µm square aperture). The observations were reduced using the ST11 interactive data processing programs developed at the Meudon observatory. The resulting redshifts are listed in table I referenced as CA1 and CA2 for the first and the second run respectively. The galaxy number is from Dressler (1980).

2.2 CALAR AUTO DATA REDUCTION. — Wavelength calibration was carried out using the Fe-Ne comparison source and night sky lines on each spectrum. Usually these set of lines are used in a fifth order polynomial fit of the wavelength solution with typical r.m.s. residuals of 0.6 Å.

The redshift computations were made using the H and K lines, the CaI 4226.70 Å one, the G band, H_γ, H_β, H_α, Mg 1b triplet and Na 1D doublet, and all these lines in the best case. Note that the [OII] 3727.30 Å line was not observed because of its position at the extreme edge of the spectral range.

Each line was fitted using a Gaussian profile so that a set of redshift was obtained depending upon the number of useful lines for each spectrum. The final redshift was determined following Balkowski *et al.* (1981), the principles of which are in de Vaucouleurs and de Vaucouleurs (1967). In order to check the accuracy of the procedure, we have observed three known galaxies in the Coma cluster (Tab. I): the results are in a very good agreement with previous values. The centroid determinations of the features measured is more or less uncertain, depending on the noise in the spectrum. This source of error is reflected for each galaxy by the dispersion of redshifts from individual lines. This dispersion is noted in table I.

2.3 OBSERVATIONS AT LA PALMA. — Observations were made during three nights (two nights were suitable for observations because of technical and clouds problems) with the 2.5 m Isaac Newton telescope in July 1984.

We used the Intermediate Dispersion Spectrograph (IDS) with a dispersion of 66.5 Å/mm. The spectral region covered was from 3500 to 5600 Å with a 632 line/mm grating blazed at 4500 Å - two slit apertures were used: 180 and 360 µm. The Image Photon Counting System (IPCS) (Boksenberg, 1982) was used as the detector with the 235 mm camera. The spectral resolution

was 2 Å FWHM. The exposure time was 1000 s so that the average signal to noise ratio measured was 15. For each galaxy, a comparison spectrum was made using a Cu-Ar source before and after each spectrum. In order to use a cross-correlation procedure for data reduction, we have taken every night two spectra, one of the G8III giant HD210807 ($V_r = -14.8 \pm 0.5$ km/s) and one of the G9III giant HD213009 ($V_r = -39.7 \pm 0.6$ km/s) using a 55 µm slit. In order to check possible mechanical flexure and instabilities the zero-points of the two stellar spectra were checked against each other, and agreed to within 20 km/s.

2.4 IPCS REDUCTIONS. — Reduction of the IPCS spectra followed the SPICA standard procedure similar to that used by many other observers (e.g. Davies, 1981; Sadler, 1982). After correction of distortion, each spectrum was flat-fielded using a normalized flat field. The comparison spectra taken immediately before and after each galaxy observation were checked for a shift and then added together. No significant shift was found.

Wavelength calibration was performed by fitting a fifth order polynomial to each arc spectrum to a wavelength scale and checking the position of several lines. Each galaxy spectrum was rebinned to a log λ scale using the corresponding calibration polynomial, then sky subtracted and normalized. After filtering to remove noise and pixel to pixel sensitivity variations the spectrum of a given galaxy is then cross-correlated with the two stars above mentioned, the measured shift $\Delta \log \lambda$ being easily converted to a velocity shift by:

$$v = c (e^{\Delta \log \lambda} - 1).$$

In order to estimate the accuracy of the radial velocities measured in this observing program, we have fit a Gaussian profile to the correlation peak displayed after a cross correlation procedure. The accuracy of the redshift determination is obtained from the width of the half height of the peak. Because of the response of the IPCS in the blue region (as for IDS at ESO) we cannot provide information on the [OII] 3727.30 Å line, specially for galaxies having H_β and H_α in emission.

2.5 OBSERVATIONS AT ESO. — Observations were made during five nights with the 1.52 m telescope at La Silla in november 1984. We used the Image Dissector Scanner (IDS) mounted on a Boller & Chivens spectrograph which ends with an output phosphor readed by a scanning device dividing the one-dimensional image into 2048 pixels. The dispersion was 114 Å/mm so that the spectral region covered was from 3830 to 6150 Å with a 600 line/mm grating blazed at 5000 Å. The slit width was 200 µm widened and the spectral resolution was about 6 Å at FWHM. The average exposure time was 3600 s so that the measured signal to noise ratio was ranging around 12. For each galaxy, a comparison was made before and after each spectrum using an He-Ar source. In order to obtain the redshift by cross-correlation procedure, we have also taken every night two spectra, one of the G3IV star HD6655 ($V_r = 15.5 \pm 0.5$ km/s)

and one of the KIII giant CD-43°2527 ($V_r = 13.1 \pm 0.3$ km/s); both stars are classical IAU Standards.

2.6 ESO DATA REDUCTION. — Data reduction was performed at Garching using the software package HIAP in order to obtain radial velocities with a cross correlation procedure. The different steps of the reduction are analogous to the IPCS ones.

2.7 OBSERVATIONS AT HAWAII. — Three galaxies have been observed at Hawaii in April 1985 with the CASS-HAWEC spectrograph installed at the Cassegrain focus of the 3.60 m telescope. The dispersion was 38 Å/min centred to observe the MgII triplet either on the galaxy and on the reference star, the giant K2III HD90861 ($V_r = 39.8 \pm 0.4$ km s⁻¹). The detector was a 236 × 236 px photon counting system with a Fe-Ne comparison source. The data reduction has been made using the cross correlation procedure, identical to the above IPCS one.

3. Estimate of errors.

In order to test the accuracy of our velocities, we have made a comparison of our redshift determinations with data available in the literature (see e.g. Palumbo *et al.*, 1983) for the same set of galaxies. The results of the comparison between the two sets of observations are summarized in table II, where we list the velocity differences between our measurements and other determinations. We have excluded three galaxies for which the discrepancy between redshift determinations was too large: A496-57, A1142-41 and A2589-17; we find that our velocities are shifted by $+4.0$ km s⁻¹ in the average with respect to the published data. The standard deviation in the difference is 89.7 km s⁻¹, which is consistent with the rms of redshift measurements mentioned in table II, ranging from 50 to 100 km/s. A valuable comparison can be made for the twelve galaxies of A496.

From the comparison of our redshifts with those of Quintana *et al.* (1985), we find that our velocities are shifted by -6.2 km s⁻¹ in the average. The standard deviation in the differences is 91.3 km s⁻¹ which is consistent with the rms uncertainty of 96.3 km s⁻¹ derived from Quintana *et al.* (1985). Note that our velocities are not as homogeneous as those measured usually, since they come from different telescopes, but all velocities in the present study agreed with those quoted by authors mentioned in table II to within ± 180 km/s (2 σ level) except for the three galaxies above mentioned, which were not included in the error calculations.

4. Conclusion.

In this paper, we have reported new redshift measurements for 100 galaxies in ten Abell clusters. These redshifts represent the initial results of the observational group on clusters of galaxies of Meudon. However, the number of redshifts collected in our survey is too much dependent of the observing time allocated and weather conditions.

The recent development of multi-object spectrographs associated with large telescopes will be a powerful tool to increase the number of redshifts collected. We are doing a major effort in this direction.

Let us emphasize that, in addition to redshifts, a lot of data like reliable magnitudes, X observations... are also required in order to improve our knowledge on clusters of galaxies and large scale structures.

Acknowledgements.

We thank the staffs of the Observatories at Calar Alto, La Palma, ESO and CFHT for their efficient help during the observing runs.

References

- BALKOWSKI, C., LE DENMAT, G., NOTTALF, L.: 1981, *Astron. Astrophys. Suppl. Ser.* **43**, 121.
 BOKSENBURG, A.: 1982, *Philos. Trans. R. Soc. London, Ser. A*: **307**, 531.
 DAVIES, R. L.: 1981, *Mon. Not. R. Astron. Soc.* **194**, 879.
 DEKEL, A.: 1984, *Astrophys. J.* **284**, 445.
 DRESSLER, A.: 1980, *Astrophys. J. Suppl. Ser.* **42**, 565.
 FADER, S. M., DRESSLER, A.: 1977, *Astron. J.* **82**, 187.
 FONTANELLI, P.: 1984, *Astron. Astrophys.* **138**, 85.
 GELLER, M. J., BEERS, T. C., BOTHUN, G. D., HUCHIRA, J. P.: 1984, *Astron. J.* **89**, 319.
 GREGORY, S. A., THOMPSON, L. A.: 1984, *Astrophys. J.* **286**, 422.
 HINZEN, P.: 1980, *Astron. J.* **85**, 626.
 HUMASON, M. L., MAYALL, N. U., SANDAGE, A. R.: 1956, *Astron. J.* **61**, 97.
 KENT, S. M., GUNN, J. E.: 1982, *Astron. J.* **87**, 945.
 MAZURE, A., GERBAL, D., PROUST, D., CAPELATO, H. V.: 1985, *Astron. Astrophys.*, in press.
 PALUMBO, G. G. C., TANZELLA-NETTI, G., VETOLANI, G.: 1983, *Catalogue of radial velocities of galaxies*, Gordon and Breach (New York).
 PEBLES, P. J. E.: 1980, *The large Scale Structure of the Universe* (Princeton University Press, Princeton, USA).
 QUINTANA, H., MELNICK, J., INFANTE, L., THOMAS, B.: 1985, *Astron. J.*, in press.
 RUBIN, V. C., FORD, W. K. Jr., THONNARD, N., ROBERTS, M. S., GRAHAM, J. A.: 1976, *Astron. J.* **81**, 687.
 SADLER, E. M.: 1982, Thesis (Australian National University).
 TRUITON, K. P.: 1972, *Mon. Not. R. Astron. Soc.* **158**, 277.
 DE VAUCOULEURS, G., DE VAUCOULEURS, A.: 1967, *Astron. J.* **72**, 730.

TABLE I (continued)

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
A 2250							
57	17 07	50.4	+7.6	45	51	50	-- LPA 14445 60
A 2589							
17	23 21	26.0	+1.6	28	28	50	-- LPA 12200 60
21	23 21	27.0	+1.0	29	01	E	-- " 12126 60
28	23 21	45.1	+1.0	30	36	S	-- " 12406 60
30	23 21	32.9	+1.6	30	21	50	-- " 12476 60
31	23 21	29.7	+1.6	32	46	E	-- " 12620 60
32	23 21	28.7	+1.6	32	13	50	-- " 12651 60
33	23 21	26.0	+1.0	31	32	E	-- " 12320 60
35	23 21	17.8	+1.6	29	36	50/A	-- " 14031 60
45	23 21	29.0	+1.0	35	45	E	-- " 12009 60
46	23 21	26.5	+1.0	35	59	E	-- " 12672 60

Notes to table I:

Column 1: Galaxy number in the cluster.

Column 2 and 3: Right ascension and declination (1950).

Column 4: Morphological type.

Column 5: Estimated total apparent visual magnitude.

Column 6: Observatory. CA1 and CA2 are observations made at Calar Alto in 1984 and 1985 respectively, LPA at La Palma in 1984, ESO at La Silla in 1984 and CF11 at Mauna Kea in 1985.

Column 7: Galaxy heliocentric velocity and estimated error.

Column 8: Notes: 1- Results are derived from two correlation peaks. 2- H α and H β in emission. 3- 3C 306.

TABLE II. — Comparison with published redshifts.

GALAXY	V-V(REF)	REFERENCE
A 151 - 61	+5	HUMASON ET AL. (1956)
62	-77	"
A 496 - 13	+58	QUINTANA ET AL. (1985)
25	+26	"
20	+911	"
32	+98	"
35	-4	"
34	-13	"
42	-93	"
46	-55	"
50	+21	"
57	+591A	"
59	+157	"
75	-140	"
77	-152	"
A 1142 - 19	-135	GILLER ET AL. (1964)
20	-90	"
29	-55	"
37	+71	"
41	-571A	"
A 1656 - 65	+143	KENT AND HORN (1962)
105	-47	"
179	+611	"
A 1983 - 15	+117	TRILLION (1972)
114	-67	HORN ET AL. (1970)
A 2063 - 60	+156	HUMASON ET AL. (1956)
A 2256 - 37	+53	FABER AND WESSLER (1977)
A 2589 - 17	+1045*	HINTZB (1980)
28	+13	"
32	+23	"

<V-V(REF)>	+4.0	
σ	+69.7	

* NOT INCLUDED IN THE CALCULATIONS