

## Phase Transitions in Coupled XY-Ising Systems

Enzo Granato,<sup>(a)</sup> J. M. Kosterlitz, and Jooyoung Lee

*Department of Physics, Brown University, Providence, Rhode Island 02912*

M. P. Nightingale

*Department of Physics, University of Rhode Island, Kingston, Rhode Island 02881*

(Received 24 October 1990)

We study the critical behavior of fully frustrated XY and Josephson-junction systems by means of a coupled XY-Ising model. From Monte Carlo and transfer-matrix calculations, we find separate XY and Ising and first-order transitions, depending on the parameters. In addition, a line of continuous transitions is found, with simultaneous loss of XY and Ising order and novel critical behavior. This result is supported by Monte Carlo simulations of frustrated XY models on square and triangular lattices.

PACS numbers: 75.40.Mg, 64.60.Cn

Models containing both continuous U(1) and discrete  $Z_2$  symmetry can give rise to interesting and unusual critical behavior. In general, one expects one of three scenarios: Transitions of the Ising and XY types occur at different temperatures or at the same temperature in a decoupled fashion, or else, in the case of strongly coupled excitations, a single transition in a different universality class. The nature of this single transition is one of the most important issues but is also less amenable to numerical or analytical approaches. A prototype of this system is a fully frustrated (FF) XY model on a triangular or a square lattice. This system has attracted great attention in recent years,<sup>1-15</sup> as it can be physically realized as a Josephson-junction array of large capacitance in a perpendicular magnetic field corresponding to a half-flux quantum per plaquette. The model is defined by the Hamiltonian

$$H/kT = - \sum_{ij} J_{ij} \cos(\theta_i - \theta_j), \quad (1)$$

where  $J_{ij} = \pm J$  ( $J > 0$ ) for  $i, j$  nearest neighbors subject to the constraint that the gauge-invariant product of  $J_{ij}$  around a plaquette is negative. For the square lattice, this can be accomplished by ferromagnetic horizontal rows and alternating ferromagnetic and antiferromagnetic columns and for the triangular lattice by isotropic antiferromagnetic couplings. This leads one to frustration and a double degeneracy in addition to the degeneracy due to the continuous symmetry.

From renormalization-group ideas and universality, it is quite natural to expect that the critical behavior of the FF XY model, arising from the interplay between XY and Ising-like excitations, could be described by a coupled XY-Ising model. In fact, a Ginzburg-Landau free energy of the FF XY models<sup>4,6-8</sup> and renormalization-group analysis leads one to consider a coupled XY-Ising model of the form<sup>8,15</sup>

$$\frac{H}{kT} = - \sum_{\langle ij \rangle} [(A + B\sigma_i\sigma_j)\cos(\theta_i - \theta_j) + C\sigma_i\sigma_j], \quad (2)$$

where  $A$ ,  $B$ , and  $C$  are effective couplings depending on the initial values of the parameters in the Ginzburg-Landau free energy.<sup>16</sup> As a consequence, the square and triangular FF XY models<sup>17</sup> can be considered to be at different initial points in the parameter space of the same model of Eq. (2). The isotropic FF XY models correspond to  $A=B$ , but a generalized version of the FF XY model on a square lattice,<sup>9</sup> where the antiferromagnetic couplings have different magnitudes compared to the ferromagnetic ones, corresponds to  $A \neq B$ .<sup>7</sup> An early study of this model<sup>8</sup> revealed a bifurcation point close to  $C=0$  in the subspace  $A=B$ . For larger positive  $C$  there is a double transition with an XY followed by an Ising as temperature is increased, while if  $C < 0$  there is a single transition with simultaneous loss of XY and Ising order. An identical topology of the phase diagram has also been found in a generalized Coulomb-gas representation of the FF XY model containing fractional charges.<sup>10</sup> An important feature of the phase structure when  $A=B$  is that Ising disorder induces XY disorder. In the model of Eq. (2) this can be seen as arising from the special coupling between the variables since a domain wall in the Ising variables leaves the XY spins uncoupled as  $1 + \sigma_i\sigma_j = 0$  at a wall. Although physically different, this is similar to the mechanism for a single transition in the FF XY model.<sup>3</sup> The role played by the fractional corner charges in that model, which, when the walls melt, trigger an unbinding of integer charges destroying XY order, is played here by the domain wall itself. Our claim is that, in spite of the apparent discrepancies between the FF XY models<sup>1,2</sup> of Eq. (1) and the Coulomb-gas representation<sup>10,11</sup> of frustrated junction arrays, both can be accommodated in the  $A=B$  subspace of Eq. (2). Changing the temperature in either corresponds to different paths through the same phase diagram. In experiments, positional disorder can lead to a double transition if there is a single one in the ideal system. These transitions, however, will be in different universality classes, as this kind of disorder acts as a ran-

dom bond for the Ising and random dipoles for integer charges in the Coulomb-gas representation.<sup>18</sup>

In this work we report the results of a detailed study by Monte Carlo (MC) simulations of the critical behavior of the coupled  $XY$ -Ising model in Eq. (2) along the line of single transitions. We have also performed MC simulations of the original FF  $XY$  models of Eq. (1) to compare the critical exponents. The phase diagram obtained by MC simulation is indicated in Fig. 1. We monitor only the Ising variables and find a segment of continuous non-Ising phase transitions starting at  $P$  which eventually becomes first order for  $C \ll 0$  at a tricritical point  $T$ . The results suggest a nonuniversal behavior as the exponents associated with the Ising order parameter vary systematically along the line  $PT$ . In addition, an evaluation of the central charge  $c$  using a Monte Carlo transfer matrix<sup>19</sup> gives a rather surprising result:  $c$  is found to vary from  $c \approx 1.5$  near  $P$  to  $c \approx 2$  near  $T$ . For the FF  $XY$  model on a square and a triangular lattice, using the same methods to evaluate the critical exponents, we find these exponents to be in fair agreement with the corresponding ones obtained for the coupled  $XY$ -Ising model near  $P$ . The central charge of the FF  $XY$  model evaluated recently by Thijssen and Knops,<sup>13</sup>  $c = 1.66(4)$ , is consistent with the corresponding result for the coupled  $XY$ -Ising model near the bifurcation point.

The simulations were performed employing the standard Metropolis algorithm using small system sizes  $L$  to achieve good statistics. Critical exponents and the location of first-order transitions were obtained from long simulations, typically of  $5 \times 10^6$  MC steps, by means




FIG. 1. Phase diagram obtained from MC simulations. Lines through the data points ( $\times$ ) are guides to the eyes. Solid and dotted lines indicate continuous and first-order transitions. Arrows indicate the known critical couplings for the Ising and  $XY$  models. Inset on left: Finite-size scaling of free-energy barrier  $\Delta F_E$  at a first-order point.

of the histogram method<sup>20</sup> and the finite-size scaling analysis of Lee and Kosterlitz (LK).<sup>21</sup> We also compared our results with those derived by applying the cumulant method<sup>22</sup> and found fair agreement.

The LK method is particularly suited to studying temperature and field-driven first-order transitions by a finite-size scaling analysis of the barriers between different phases at the transition. At a temperature-driven first-order transition, the free-energy barrier  $\Delta F_E$  between ordered and disordered states is obtained from the energy histogram  $N(E; A, C, L)$  as the height of the peak in  $-\ln N(E)$ . At a first-order transition  $\Delta F_E$  is a monotonically increasing function of  $L$  and we show the results for  $A=6.5$  in an inset of Fig. 1. As we move along the transition line towards  $P$ , this peak becomes lower and disappears at  $T$ . We identify this as a tricritical point but this may overestimate the range of continuous transitions since the absence of a peak in systems of limited size does not exclude a very weak first-order transition.

Along the segment  $TP$ , we assume that the transition is continuous and, in principle, there are two sets of exponents to be determined since there are both Ising and  $XY$  variables. We are able to estimate only those associated with the former to any degree of accuracy and they are obtained from the histogram of Ising magnetization  $m$ . The corresponding free-energy barrier  $\Delta F_m$  between the bulk states  $\pm m$  increases as  $L^{1/\nu}$  for  $L < \xi$  in the ordered state and decreases to zero in the disordered state. This change of behavior can be used to accurately determine the critical values of  $A$  and  $C$ . More importantly, the exponent  $\nu$  can be determined from  $S = \partial \Delta F_m / \partial C \approx L^{1/\nu}$  without a precise determination of the critical parameters. This is a strength of the LK method as it yields  $\nu$  from a one-parameter fit<sup>23</sup> in contrast to most earlier MC analyses<sup>1,2</sup> which involve fitting at least two. We identify  $\nu \neq 1$  as a single transition since, if the Ising and  $XY$  transitions are separate or decoupled,  $\nu=1$  at the Ising transition. We were unable to obtain meaningful results for the helicity modulus near the transition so the line  $XP$  in Fig. 1 is an interpolation from the known position in the phase with  $m=1$ . We identify  $P$  as the bifurcation point but cannot exclude the possibility of a short segment of single decoupled transition with  $\nu=1$  to the left of  $P$ . The exponent  $2\beta/\nu$  is obtained from the  $L$  dependence of the peak separation in the histogram of  $m$  which scales as  $L^{-\beta/\nu}$  at criticality, although this is subject to larger uncertainties than  $\nu$ . Our estimates of the Ising exponents  $\nu$  and  $2\beta/\nu$  are plotted in Fig. 2 along the line  $PT$ . It is clear that these differ from the pure Ising values and seem to vary systematically on the transition line indicating nonuniversal behavior. We have also made an independent study using the cumulant method<sup>22</sup> and find consistency.<sup>15</sup>

In order to compare the result for the critical ex-




FIG. 2. Critical exponents obtained from a finite-size scaling of  $\Delta F_m$  on lattices of size  $10 \leq L \leq 30$ . Horizontal arrows correspond to known results and the vertical arrow indicates the estimated location of the bifurcation point in Fig. 1.

ponents of the coupled  $XY$ -Ising model near the bifurcation point with those of the FF  $XY$  model, we used the same methods to evaluate the Ising-like critical exponents for the triangular and the square lattices.<sup>24</sup> Figure 3 shows the result of a finite-size scaling analysis of  $S = \partial \Delta F_\chi / \partial J$  for both cases.  $\Delta F_\chi$  was obtained from histograms of the staggered chirality order parameter  $\chi$ .<sup>2,3</sup> A small curvature is still observed even for the largest system sizes, indicating that correction to scaling is still important. However, if one estimates the exponents from the largest sizes, one obtains  $\nu = 0.83(4)$ ,  $2\beta/\nu = 0.28(4)$  (triangular) and  $\nu = 0.85(3)$ ,  $2\beta/\nu = 0.31(3)$  (square). Considering the curvature in the data, these can be considered to be upper bounds to the asymptotic result. Even so, they do differ significantly from the pure Ising exponents and seem to be consistent with the values obtained near the bifurcation point in Fig. 1. Note that the correlation lengths probed in the simulations of the FF  $XY$  models are much smaller than the ones in the coupled  $XY$ -Ising model since the size of the unit cell is larger. Although the exponents for the triangular and the square lattices also seem to agree with each other within the estimated uncertainties, it is perfectly possible that they may eventually differ if simulations are carried out in sufficiently large systems. This nonuniversal behavior of the FF  $XY$  model would be in complete agreement with our findings for the coupled  $XY$ -Ising models.

Additional information on the nature of the critical behavior on the line  $PT$  is provided by the central charge  $c$  which, to be consistent with varying exponents,<sup>25</sup> must obey  $c \geq 1$ . Preliminary investigations, using MC




FIG. 3. Finite-size scaling of the peak position  $\chi^2$  and  $S = \partial \Delta F_\chi / \partial J$  from the histogram of the chirality  $\chi$  for the square ( $\square$ ) and the triangular ( $\Delta$ ) FF  $XY$  models. Solid lines are least-squares fits to  $L > 10$  data.

transfer-matrix methods<sup>19</sup> on infinite strips of width up to  $L = 16$ , yield the very surprising result that  $c$  seems to vary continuously from  $c \approx 1.5$  at  $P$  to  $c \approx 2$  at  $T$ . This implies that the line  $PT$  cannot be described as a superposition of critical  $XY$  and Ising lines with  $c = 1.5$ , as suggested by Foda,<sup>26</sup> but this is a completely new effect. The only models where  $c$  is known to vary are models that do not have a symmetric transfer matrix, e.g., the  $q$ -state Potts and  $O(n)$  models with a continuously varying number of states  $q$  and  $n$ . Unlike in those models, on the line  $PT$  a parameter is changing that does not affect the symmetry. It seems to us that there are three possible explanations: (i) The system is not conformally invariant on the line  $PT$ ; (ii) the result is an artifact of limited strip widths; and (iii) it is a real and interesting effect. Further investigation is under way.

We note that this variation of  $c$  is entirely consistent with the Zamolodchikov  $c$  theorem<sup>27</sup> which states that there exists a quantity  $c(l)$  such that  $dc(l)/dl \leq 0$  along a renormalization-group trajectory and tends to  $c$  at a fixed point. The model of Eq. (2) may be obtained by adding a relevant term  $h_2 \cos 2(\theta_i - \phi_i)$  to two coupled  $XY$  models<sup>7</sup> which implies  $c \leq 2$ . If, on the other hand, we start from decoupled critical  $XY$  and Ising models and add a term  $B \sigma_i \sigma_j [1 - \cos(\theta_i - \theta_j)]$ , we obtain  $c \geq \frac{3}{2}$  since such a term is irrelevant about the decoupled fixed point for  $B$  small.<sup>7,28</sup> Since the true model is at  $A = B$ , this yields the allowed range of  $\frac{3}{2} \leq c \leq 2$ . We finally note that the value of  $c = 1.66$  of Thijssen and Knops<sup>13</sup> for the FF  $XY$  model using the same methods is within the allowed range.

We thank R. M. Marinho, Jr., R. H. Swendsen, K.

Binder, T. Ala-Nissila, and H. W. J. Blöte for helpful discussions. This work was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (E.G.) and NSF Grants No. DMR-8918358 (J.M.K. and J.L.) and No. DMR-8704730 (M.P.N.). Two of us (E.G. and J.M.K.) acknowledge Fundação de Amparo e Pesquisa do Estado de São Paulo (FAPESP) (Brazil) for supporting a visit (J.M.K.) to the Instituto de Pesquisas Espaciais where part of this work was done.

<sup>(a)</sup>On leave from Instituto Nacional de Pesquisas Espaciais, 12201 São José dos Campos, São Paulo, Brazil.

<sup>1</sup>J. Villain, *J. Phys. C* **10**, 4793 (1977); S. Teitel and C. Jayaprakash, *Phys. Rev. B* **27**, 598 (1983); *Phys. Rev. Lett.* **51**, 199 (1983); W. Y. Shih and D. Stroud, *Phys. Rev. B* **32**, 158 (1985).

<sup>2</sup>S. Miyashita and H. Shiba, *J. Phys. Soc. Jpn.* **53**, 1145 (1984); D. H. Lee, J. D. Joannopoulos, J. W. Negele, and D. P. Landau, *Phys. Rev. B* **33**, 450 (1986); E. van Himbergen, *Phys. Rev. B* **33**, 7857 (1986).

<sup>3</sup>T. C. Halsey, *J. Phys. C* **18**, 2437 (1985); S. E. Korshunov and G. V. Uimin, *J. Stat. Phys.* **43**, 1 (1986).

<sup>4</sup>M. Y. Choi and S. Doniach, *Phys. Rev. B* **31**, 4516 (1985); M. Y. Choi and D. Stroud, *Phys. Rev. B* **32**, 5773 (1985).

<sup>5</sup>P. Minnhagen, *Phys. Rev. B* **32**, 3088 (1985).

<sup>6</sup>M. Yosefin and E. Domany, *Phys. Rev. B* **32**, 1778 (1985).

<sup>7</sup>E. Granato, J. M. Kosterlitz, and J. Poulter, *Phys. Rev. B* **33**, 4767 (1986); E. Granato and J. M. Kosterlitz, *J. Phys. C* **19**, L59 (1986); *J. Appl. Phys.* **64**, 5636 (1988).

<sup>8</sup>E. Granato, *J. Phys. C* **20**, L215 (1987).

<sup>9</sup>B. Berge, H. T. Diep, A. Ghazali, and P. Lallemand, *Phys. Rev. B* **34**, 3177 (1986).

<sup>10</sup>J. M. Thijssen and H. J. F. Knops, *Phys. Rev. B* **37**, 7738

(1988); J. M. Thijssen, *Phys. Rev. B* **40**, 5211 (1989); H. Eikmans, J. E. van Himbergen, H. J. F. Knops, and J. M. Thijssen, *Phys. Rev. B* **39**, 11759 (1989).

<sup>11</sup>G. Grest, *Phys. Rev. B* **39**, 9267 (1989).

<sup>12</sup>M. Gabay, T. Garel, G. N. Parker, and W. M. Saslow, *Phys. Rev. B* **40**, 264 (1989).

<sup>13</sup>J. M. Thijssen and H. J. F. Knops, *Phys. Rev. B* **42**, 2438 (1990).

<sup>14</sup>H. Kawamura, *Phys. Rev. B* **38**, 960 (1988).

<sup>15</sup>J. Lee, E. Granato, and J. M. Kosterlitz (unpublished).

<sup>16</sup>The same model also describes the  $T=0$  superconductor-insulator transition in a Josephson-junction ladder at half-flux quantum; E. Granato, *Phys. Rev. B* **42**, 4797 (1990).

<sup>17</sup>For triangular FF XY models, Ref. 4 obtains a free energy with coupling between the phases only in sixth order. A change of variables, however, leads to a free energy which to fourth order has the same form as the square case, Refs. 14 and 15.

<sup>18</sup>E. Granato and J. M. Kosterlitz, *Phys. Rev. Lett.* **62**, 823 (1989).

<sup>19</sup>M. P. Nightingale and H. W. J. Blöte, *Phys. Rev. Lett.* **60**, 1562 (1988).

<sup>20</sup>A. M. Ferrenberg and R. H. Swendsen, *Phys. Rev. Lett.* **61**, 2635 (1988); **63**, 1195 (1989); M. Falcioni, E. Marinari, M. L. Paciello, G. Parisi, and B. Taglienti, *Phys. Lett.* **108B**, 331 (1982); E. Marinari, *Nucl. Phys.* **B235**, 123 (1984).

<sup>21</sup>J. Lee and J. M. Kosterlitz, *Phys. Rev. Lett.* **65**, 137 (1990); *Phys. Rev. B* **43**, 3265 (1991).

<sup>22</sup>K. Binder, *Phys. Rev. Lett.* **47**, 693 (1981).

<sup>23</sup>H. Park and M. den Nijs, *J. Phys. A* **22**, 3663 (1989); H. Park, *J. Phys. A* **23**, 1789 (1990).

<sup>24</sup>J. Lee, J. M. Kosterlitz, and E. Granato (unpublished).

<sup>25</sup>J. L. Cardy, *J. Phys. A* **20**, L891 (1987).

<sup>26</sup>O. Foda, *Nucl. Phys.* **B300**, 611 (1988).

<sup>27</sup>A. B. Zamolodchikov, *Pis'ma Zh. Eksp. Teor. Fiz.* **43**, 567 (1986) [*JETP Lett.* **43**, 733 (1986)].

<sup>28</sup>M. den Nijs, *Phys. Rev. B* **32**, 4785 (1985).