



Special Edition

Data assimilation for nowcasting in the terminal area of Janeiro

Assimilação de dados para previsão na área terminal do Rio de Janeiro

Vinícius Albuquerque de Almeida ^I

Gutemberg Borges França ^{II}

Haroldo Fraga de Campos Velho ^{III}

^I Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil. E-mail: vinicius@lma.ufrj.br.

^{II} Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil. E-mail: gutemberg@lma.ufrj.br.

^{III} Instituto Nacional de Pesquisas Espaciais, São Paulo, Brazil. E-mail: haroldo.camposvelho@inpe.br.

ABSTRACT

The process of data assimilation, in which meteorological observations and weather forecasts are merged to provide an analysis field, has been largely studied by the scientific and operational centers. The 3D-Variational (3D-Var) approach available in the Weather Research and Forecast (WRF) computer model is evaluated for data assimilation for the Terminal Control Area of Rio de Janeiro (TCA-RJ). The basic goal of any variational data assimilation system is to provide an optimal estimate of the atmospheric state at analysis time. The analysis field is estimated by combining a first guess (previous forecast) and an observation field, weighted by the error matrices. This system was designed for nowcasting (forecasts up to 6h) for the TCA-RJ through assimilation of surface, sounding, and wind profile data. The preliminary results show the model sensitivity to each observation type and encourage the use of this technique operationally for the support of air traffic management controlled by the Brazilian Air Force.

Keywords: Data assimilation; WRF; 3D-Var; Surface data; Profile data.

RESUMO

O processo de assimilação de dados, onde observações meteorológicas e previsões do tempo são combinadas para fornecer um campo de análise, tem sido muito estudado pela comunidade científica e operacional.

científica e por centros operacionais. A utilização do método variacional 3D (3D-Var) implementado no modelo Weather Research and Forecast (WRF) é avaliada para assimilação de dados do Aeroporto Internacional do Rio de Janeiro (TCA-RJ). O objetivo principal de qualquer método variacional de assimilação de dados é produzir uma estimativa ótima do estado atmosférico no momento da análise. O campo de análise é estimado por um campo de estimativa inicial (campo de dados anterior) e um campo de observações, ponderados pelas matrizes de erro. O WRF é utilizado para nowcasting (previsão de até 6h) para a TCA-RJ através de ciclos de assimilação de dados de superfície, sondagem e perfilhadores de vento. Os resultados preliminares sobre a sensibilidade de cada tipo de observação e encorajam a utilização desta técnica operacional para fornecer suporte ao controle de tráfego aéreo controlado pela Força Aérea Brasileira.

Palavras-chave: Assimilação de dados; WRF; 3D-Var; Dados de superfície; Dados de perfil

1 INTRODUCTION

Numerical weather prediction (NWP) is considered an initial-value problem, where the current state of the atmosphere is used as input to a numerical model for simulating or predicting its evolution over space and time. The problem of determination of the initial conditions for the model is very important and complex, and has become a science in itself (DALEY, 1999). Various methods have been developed since the 1950s to tackle this problem. Daley (1991) and Daley (2012) can be used to a broader review on data analysis and assimilation techniques.

In meteorology, there is a wide variety of data sources able to be assimilated to estimate the state of the atmosphere, including conventional and non-conventional data. Conventional data include surface observations, balloon soundings, aircraft and ship observations. On the other hand, data retrieved from satellites (e.g. radiance), wind profilers (e.g. LiDAR), and radar are usually known as non-conventional. Conventional data are usually assimilated in global models, but such data very often represent local conditions and are smoothed due to interpolation methods and quality control routines. Also, some observations are not from the observation network, and they are not processed by data assimilation procedures used in the global models. Therefore, the best determination for the atmosphere state for a resolution limited area model is obtained by assimilating local retrieved data.

Preliminary tests on the sensibility of a regional model are evaluated for the assimilation of profile data (sounding) in the terminal control area of Rio de Janeiro (Brazil).

2 MATERIAL AND METHODS

2.1 Data and study area

The experiments performed in this work used (i) data from Global Forecast System (GFS) forecasts for model initialization and boundary conditions; (ii) METAR/SPECI data retrieved from the surface meteorological stations at several airports in Rio de Janeiro (Figure 1).

sounding data retrieved daily at the International Airport of Rio de Janeiro, known as Gal (SBGL). Table 1 show details of the used data and its source for reproducibility. Figure 2 study domain and also the model grid used on the simulation runs.


Table 1 – Source, description, and frequency of the data used for assimilation sensibility

Source	Description	Frequency	Da
GFS https://rda.ucar.edu/datasets/ds084.1	Initial and boundary atmospheric conditions at 0.25-degree resolution	3 h	2
METAR/SPECI data https://www.redemet.aer.mil.br	Surface data at airports	1 h	1
Atmospheric Sounding https://www.redemet.aer.mil.br	Atmospheric profiles of SBGL of temperature, relative humidity, atmospheric pressure, winds and sounding-derived atmospheric instability indices.	daily (0 and 12Z)	1

2.2 WRF model

The Weather Research and Forecasting (WRF) Model is a next-generation numerical weather prediction system designed for both atmospheric research and forecasting applications. The code embraces two dynamical cores, a data assimilation system, a software architecture supporting parallel computation and system extensibility. The develop WRF began in the latter 1990's, and was a collaborative partnership of the National Center for Atmospheric Research (NCAR), the National Oceanic and Atmospheric Administration (represented by the National Centers for Environmental Prediction (NCEP) and the Earth System Research Laboratory), the U.S. Air Force, the Naval Research Laboratory, the University of Oklahoma, and the Federal Aviation Administration (FAA). Please refer to the WRF User's Guide and the Technical Note document available at <http://www2.mmm.ucar.edu/wrf/> for the completeness of the 3D-Var implementation present at WRF (SKAMAROCK et al., 2019)

Figure 1 – Terrain map and airport locations



The WRF model solves a set of equations for atmospheric state evolution, which include (i) conservation of momentum; (ii) thermodynamic energy conservation; (iii) mass conservation; (iv) the geopotential relation; and (v) the equation of state. Also, several physical processes are parameterized, because they may be too small, too brief, too complex, too poorly understood, or too computationally costly to be explicitly represented, e.g. short and longwave radiative transfer, planetary boundary layer, turbulence, cumulus convection, cloud microphysics, and precipitation.

The WRF model was integrated in a 5-km grid with 35 levels in vertical, generating outputs from surface and pressure-level variables. Regarding the parametrizations, the options were chosen: Microphysics - WRF Single-moment 3 (HONG *et al.*, 2004), Grell-Freitas Ensemble Scheme (GRELL AND FREITAS, 2014), Radiation - Dudhia Scheme (DUDHIA, 1989) / RRTM Longwave Scheme (MLAWER *et al.*, 1997), Planetary Boundary Layer - Yonsei University Scheme (YSU) (HONG, 2006), and Land-Surface model: Urban Canopy Model (TEWARI *et al.*, 2004).

2.3 3D-Var

The 3D-Var approach was selected, as implemented in the Data Assimilation component of the WRF framework. An introduction to the basic ideas of variational data assimilation is specifically the WRF Data Assimilation (WRFDA) system is deeply discussed in Barker (2004) and Barker *et al.* (2012).

Among various data assimilation methods, the variational approaches have been widely used in meteorology, specifically the method 3D-Var. In the 3D-Var approach, a cost function (equation 1) is defined as the difference between observation (\vec{y}^o) and the analysis value (\vec{x}) at the observational grid $[H(\vec{x})]$ under norm-R, regularized by the difference between the analysis value and the background (\vec{x}^b) under norm-B (SASAKI, 1970; KALNAY, 2012). The analysis field is obtained by the direct minimization of such function. Matrices for both, the background (R) and covariance (B) are required.

(B), are considered in the minimization process. The operator H transforms the gridded the observation space for comparison against the observation vector \vec{y}^o .

$$J = \frac{1}{2} \{ [\vec{y}^o - H(\vec{x})]^T R^{-1} [\vec{y}^o - H(\vec{x})] + (\vec{x} - \vec{x}^b)^T B^{-1} (\vec{x} - \vec{x}^b) \}$$

In essence, the 3D-Var approach consists in processing observed information in window (typically from 1 h before the analysis time to 1 h after) over a spatial domain process, a subset of the observed data is retrieved to be assimilated in a previous forec minimizing the cost function.

2.4 Experiment


The WRF model was configured on the spatial domain shown in figure 2, in a resolution. The domain is centered in the metropolitan area of Rio de Janeiro and encompasses the airports described in figure 1, where the observations are retrieved.

The period of 72 hours was considered for the sensibility test, from Jan 1st 0Z to Jan 4th 0Z, 2017. This period was randomly chosen for the preliminary results, but a larger period (several years) will be considered in the future.

The initial and boundary conditions for the control run were obtained from the degree resolution and consists of gridded meteorological data for the study period.

The experiment was performed in three steps: (i) a control run was performed for Jan 1st 0Z to Jan 4th 0Z, 2017; (ii) a synthetic observation run, where Gaussian white noise was added to the GFS analysis field at SBGL location, and then the model was integrated without data assimilation; and (iii) an assimilation run where a synthetic profile of temperature and dew-point temperature, extracted from the observation run at SBGL location, was assimilated at the initial timestep.

Figure 2 – Horizontal computational domain used for model executions



3 RESULTS AND DISCUSSION

This section presents the results of the experiments performed in this work and the spatial-temporal characteristics of data assimilation on study area.


Data assimilation result for air temperature difference at 850 hPa is shown in Figure 3a, and the difference for the initial condition (Jan 1st, 2017 0Z) between experiment-(ii) and experiment-(i) is displayed in Figure 3a, and the difference between experiment-(ii) and experiment-(iii) is in Figure 3b. The same differences are displayed in Figures 3c and 3d on Jan 1st, 2017 12Z, respectively.

The analysis field (Figures 3a and 3b) shows smoother fields from the assimilated data. The difference from the observation field mainly at the airport location and its surrounding area illustrates the smoothing. After 12 h of integration (Figures 3c and 3d), the differences between the assimilated fields, but still the assimilation field presents less difference in comparison to the observation field. The results from Figure 3 indicate that even a single temperature profile assimilation at a timestep can positively impact the analysis field.

Figure 3 – The 850-hPa air temperature field at 00Z for: (a) the difference between experiment-(ii) and experiment-(i); and (b) the difference between experiment-(ii) and experiment-(iii). At 12Z, Figures (c) and (d) show the same difference, respectively, after 12Z of integration

integration approximates to 72 h of integration, the errors increase significantly both in experiment-(i) and experiment-(iii), as shown by the blue rectangle in Figure 4. However, it is possible that the effect of the temperature profile assimilation at the initial timestep, since the errors in experiment-(iii) are smaller than the errors in experiment-(i).

Figure 4 – Vertical cross-section and time evolution at SBGL for: (a) the difference between observation field (experiment-(ii)) and experiment-(i); and (b) the difference between observation field and experiment-(iii). Red square represents the short-forecast time-window (24 h) and the blue square highlights the medium-range forecast, close to 72 h



4 CONCLUSION

The 3D-Var approach of the WRF framework was evaluated for the assimilation of temperature profile at the Galeão International Airport, Rio de Janeiro (SBGL) for a 24-h period in January 2017.

Preliminary results showed that the assimilation routine was able to adjust the temperature profile at the airport, and also keep these atmospheric characteristics present in the study domain.

The experiments performed have shown the positive impact of the assimilation on the model's overall performance. As shown by the red rectangle in Figure 4, the assimilation can be effective for the nowcasting time-window, under 24-h. Also, even after 72 h the forecast errors are smaller than no-assimilation forecasts, encouraging the use of such methods to increase reliability in those forecasts.

In the future the assimilation of different data types will be evaluated (e.g. profiler, SODAR and LiDAR) and the assimilation will be expanded for a greater period, evaluating the performance for reproducibility of different meteorological events.

ACKNOWLEDGMENTS

This study is funded by the Department of Airspace Control (DECEA) via the Organization for Scientific and Technological Development of Airspace Control (CTCEA/002-2018/COPPETEC_CTCEA). We would like to thank the National Council for Scientific and Technological Development (CNPq, Portuguese) for the research grant for the author (CNPq: 312924/2017-8) and GBF (CNPq: 304441/2018-0).

REFERENCES

- BARKER, D. M., W. HUANG, Y.-R. GUO, A. BOURGEOIS, AND X. N. XIAO. **Three-Dimensional Variational Data Assimilation System for MM5: Implementation and Results**. *Mon. Wea. Rev.*, 132, 897–914, 2004.
- BARKER, D. M., X.-Y. HUANG, Z. LIU, T. AULIGNE, X. ZHANG, S. RUGG, A. A. AL KADHI, BOURGEOIS, J. BRAY, Y. CHEN, M. DEMIRTAS, Y. GUO, T. HENDERSON, W. HUANG, LIN, J. MICHALAKES, S. RIZVI, X.-Y. ZHANG. **The Weather Research and Forecast Model's Community Variational/Ensemble Data Assimilation System: WRFDA**. *J. Meteor. Soc.*, 93, 831–843, 2012.
- CINTRA, R.S., CAMPOS VELHO, H. F. **Global Data Assimilation Using Artificial Neural Networks In Speedy Model**. *Proceedings of the 1st International Symposium on Uncertainty Quantification and Stochastic Modeling*, 2012.
- DALEY, R. **Atmospheric data analysis**. Cambridge University Press, 1991.
- DUDHIA, J. **Numerical study of convection observed during the Winter Monsoon Experiment using a mesoscale two-dimensional model**. *J. Atmos. Sci.*, 46, 3077–3107, 1989.

GRELL, G. A. AND FREITAS, S. R. **A scale and aerosol aware stochastic convection parameterization for weather and air quality modeling.** *Atmos. Chem. Phys.*, **14**, 1175–1192, 2014.

HONG, SONG-YOU, JIMY DUDHIA, AND SHU-HUA CHEN. **A revised approach to microphysical processes for the bulk parameterization of clouds and precipitation.** *Mon. Wea. Rev.*, **132**, 103–120, 2004.

HONG, SONG-YOU, YIGN NOH, JIMY DUDHIA. **A new vertical diffusion package with explicit treatment of entrainment processes.** *Mon. Wea. Rev.*, **134**, 2318–2341, 2006.

KALNAY, E. **Atmospheric Modeling, Data Assimilation and Predictability.** Cambridge University Press, 2012.

MLAWER, ELI. J., STEVEN. J. TAUBMAN, PATRICK. D. BROWN, M. J. IACONO, AND J. CLOUGH. **Radiative transfer for inhomogeneous atmospheres: RRTM, a correlated-k model for the longwave.** *J. Geophys. Res.*, **102**, 16663–16682, 1997.

SKAMAROCK, W. C., J. B. KLEMP, J. DUDHIA, D. O. GILL, Z. LIU, J. BERNER, W. W. POWERS, M. G. DUDA, D. M. BARKER, AND X.-Y. HUANG. **A Description of the NCAR Community Climate Model, Research WRF Version 4.** NCAR Tech. Note NCAR/TN-556+STR, 145 pp, 2019.

TEWARI, M., F. CHEN, W. WANG, J. DUDHIA, M. A. LEMONE, K. MITCHELL, M. EK, C. J. WEGIEL, AND R. H. CUENCA. **Implementation and verification of the unified NCEP surface model in the WRF model.** 20th conference on weather analysis and forecasting, 11–15, 2004.